# bottom

Bar lines indicate relative size

# Stream Insects & Crustaceans

### GROUP ONE TAXA

Pollution sensitive organisms found in good quality water.

- 1 Stonetly: Order Plecoptera. 1/2" 1 1/2", 6 legs with hooked tips, antennae, 2 hair-like tails. Smooth (no gills) on lower half of body. (See arrow.)
- 2 Caddistly: Order Trichoptera. Up to 1", 6 hooked legs on upper third of body, 2 hooks at back end. May be in a stick, rock or leaf case with its head sticking out. May have fluffy gill tufts on underside.
- 3 Water Penny: Order Coleoptera. 1/4", flat saucer-shaped body with a raised bump on one side and 6 tiny legs and fluffy gills on the other side. Immature beetle.
- 4 Riffle Beetle: Order Coleoptera. 1/4", oval body covered with tiny hairs, 6 legs, antennae. Walks slowly underwater. Does not swim on surface.
- 5 Mayfly: Order Ephemeroptera. 1/4" 1". brown, moving, plate-like or feathery gills on sides of lower body (see arrow), 6 large hooked legs, antennae, 2 or 3 long, hair-like tails. Tails may be webbed together.
- 6 Gilled Snail: Class Gastropoda. Shell opening covered by thin plate called operculum. When opening is facing you, shell usually opens on right.
- 7 Dobsonfly (Hellgrammite): Family Corydalidae. 3/4" 4", dark-colored, 6 legs, large pinching jaws, eight pairs feelers on lower half of body with paired cotton-like gill tufts along underside, short antennae, 2 tails and 2 pairs of hooks at back end.


### GROUP TWO TAXA

Somewhat pollution tolerant organisms can be in good or fair quality water.

- 8 Crayfish: Order Decapoda. Up to 6", 2 large claws, 8 legs, resembles small lobster.
- 9 Sawbug: Order Isopoda. 1/4" 3/4", gray oblong body wider than it is high, more than 6 legs, long antennae.

## Save Our Streams

Izaak Walton League of America 707 Conservation Lane Gaithersburg, MD 20878-2983 1(800)BUG-IWLA



Bar lines indicate relative size

### **GROUP TWO TAXA CONTINUED**

- 10 Scud: Order Amphipoda. 1/4", white to grey, body higher than it is wide, swims sideways, more than 6 legs, resembles small snrimp.
- 11 Alderfly Larva: Family Sialidae. 1" long. Looks like small hellgrammite but has 1 long, thin, branched tail at back end (no hooks). No gill tufts underneath.
- 12 Fishfly Larva: Family Corydalidae. Up to 1 1/2" long. Looks like small hellgrammite but often a lighter reddish-tan color, or with yellowish streaks. No gill tufts underneath.
- 13 Damselfly: Suborder Zygoptera. 1/2" 1". large eyes, 6 thin hooked legs, 3 broad oar-snaped tails, positioned like a tripod. Smooth (no gills) on sides of lower half of body. (See arrow.)
- 14 Watersnipe Fly Larva: Family Athericidae (Atherix). 1/4" - 1", pale to green, tapered body, many caterpillar-like legs, conical head, feathery "horns" at back end.
- 15 Crane Fly: Suborder Nemalocera. 1/3" 2", milky, green, or light brown, plump caterpillar-like segmented body, 4 finger-like lobes at back end.
- 16 Beetle Larva: Order Coleoptera. 1/4" 1", light-colored, 6 legs on upper half of body, feelers, antennae.
- 17 Dragon Fly: Suborder Anisoptera. 1/2" 2", large eyes, 6 hooked legs. Wide oval to round abdomen.
- 18 Clam: Class Bivalvia.

### **GROUP THREE TAXA**

Pollution tolerant organisms can be in any quality of water.

- 19 Aquatic Worm: Class Oligochaeta. 1/4" 2", can be very tiny; thin worm-like body.
- 20 Midge Fly Larva: Suborder Nematocera. Up to 1/4", dark head, worm-like segmented body, 2 tiny legs on each side.
- 21 Blackfly Larva: Family Simulidae. Up 1/4", one end of body wider. Black head, suction pad on other end.
- 22 Leech: Order Hirudinea. 1/4" 2", brown, slimy body, ends with suction pads.
- 23 Pouch Snail and Pond Snails: Class Gastropoda. No operculum. Breathe air. When opening is facing you, snell usually opens on lett.
- 24 Other Snails: Class Gastropoda. No operculum. Breathe air. Snail shell coils in one plane.

